

Van der Waals Density Functional for General Geometries

M. Dion,¹ H. Rydberg,² E. Schröder,² D. C. Langreth,¹ and B. I. Lundqvist²

¹Center for Materials Theory, Department of Physics and Astronomy, Rutgers University, Piscataway, NJ 08854-8019

²Department of Applied Physics, Chalmers University of Technology and Göteborg University, SE-412 96 Göteborg, Sweden

(Dated: 30 January 2004)

A scheme within density functional theory is proposed that provides a practical way to generalize to unrestricted geometries the method applied with some success to layered geometries [H. Rydberg, *et al.*, Phys. Rev. Lett. **91**, 126402 (2003)]. It includes van der Waals forces in a seamless fashion. By expansion to second order in a carefully chosen quantity contained in the long range part of the correlation functional, the nonlocal correlations are expressed in terms of a density-density interaction formula. It contains a relatively simple parametrized kernel, with parameters determined by the local density and its gradient. The proposed functional is applied to rare gas and benzene dimers, where it is shown to give a realistic description.

PACS numbers: 31.15.Ew, 71.15.Mb, 61.50.Lt

Density functional theory (DFT) for molecules and materials is widely applied with approximate *local* and *semilocal* density functionals for the interaction effects. For largely homogeneous systems, for example, simple metals and semiconductors, the local-density approximation (LDA) for these effects is appropriate. For inhomogeneous systems, for example, transition metals, ionic crystals, compound metals, surfaces, interfaces, and some chemical systems, semilocal-density approximations, like members of the family of generalized gradient approximations (GGA), work well. Today DFT describes cohesion, bonds, structures, and other properties very well for dense molecules and materials, as shown by recent studies for both single molecules [1] and dense solid-state [2] systems. However, sparse systems, including soft matter, van der Waals complexes, and biomolecules, are at least as abundant. They have interparticle separations, for which *nonlocal*, long-ranged interactions, such as van der Waals (vdW) forces, are influential.

The aim of this letter is to develop and apply a van der Waals density functional (vdW-DF) for general geometries to supplement the planar vdW-DF that we recently applied with some success [3] to several layered materials. The simplest form for the nonlocal correlation-energy part to such a functional is

$$E_c^{\text{nl}} = \frac{1}{2} \int d^3r d^3r' n(\vec{r}) \phi(\vec{r}, \vec{r}') n(\vec{r}'), \quad (1)$$

where $\phi(\vec{r}, \vec{r}')$ is some given, general function depending on $\vec{r} - \vec{r}'$ and the densities n in the vicinity of $\vec{r}$ and $\vec{r}'$. It is approximately derived here and applied to some key cases, with results that give promise for broader applications.

We start with the same approximation scheme used for layered systems [3, 4], and divide the correlation energy into two pieces,

$$E_c[n] = E_c^0[n] + E_c^{\text{nl}}[n], \quad (2)$$

which are treated in different approximations. In particular, as discussed in Ref. 4, we treat the second term

in the full potential approximation (FPA), which is exact at long distances between separated fragments, and therefore adopt Eq. (25) of Ref. 4,

$$E_c^{\text{nl}} = \int_0^\infty \frac{du}{2\pi} \text{tr} [\ln(1 - V\tilde{\chi}) - \ln \epsilon], \quad (3)$$

where $\tilde{\chi}$ is the density response to a fully self-consistent potential with long-range, interfragment spectator [5] contributions omitted [4]. V is the interelectronic Coulomb interaction, ϵ an appropriately approximated dielectric function, and u the imaginary frequency. While Eq. (3) is taken to be the definition of E_c^{nl} , we will show that Eq. (1) can be obtained with suitable approximations. The first term E_c^0 , defined by Eqs. (2) and (3), is also nonlocal; however, with the long-range vdW terms treated separately, it seems a reasonable approximation to treat E_c^0 in the LDA, and this is what we do. There is no double counting, because for a uniform system $1 - V\tilde{\chi} = \epsilon$, which implies that the LDA for E_c^{nl} vanishes. This is also the key to a seamless theory.

For layered systems the scheme was made tractable by use of the lateral average of the densities to calculate the interplanar contribution from Eq. (3). For general geometries, we make the scheme tractable by expanding Eq. (3) to second order in $S \equiv 1 - \epsilon^{-1}$, obtaining

$$E_c^{\text{nl}} \approx \int_0^\infty \frac{du}{4\pi} \text{tr} \left[S^2 - \left(\frac{\nabla S \cdot \nabla V}{4\pi e^2} \right)^2 \right]. \quad (4)$$

This vanishes in the uniform limit as it must.

To assess the nature of this approximation, we consider its prediction for the long-range interaction between two fragments. As in Ref. 6, we obtain the standard formula for the coefficient of the R^{-6} term in terms of the frequency dependent polarizability tensor $\alpha_{ij}^p(\omega)$ for each fragment $p = 1, 2$. We find that $\alpha_{ij}^p = \delta_{ij} \int d^3x \int d^3y S^p(\vec{x}, \vec{y}) / 4\pi$. We note that $S(\vec{x}, \vec{y}) / 4\pi$ is the polarization P at point $\vec{x}$ induced by the D -field at point $\vec{y}$, *i.e.*, $\vec{D}(\vec{y})$ (in this case the field from fluctuations on the other fragment). The integration over x then

gives the dipole moment, and the integration over y implies that this D -field is assumed to be spatially constant. This means that, although a local-field correction is included, it is not calculated self-consistently. It is the same approximation for the electrodynamics that was used to calculate the C_6 coefficient successfully over a large set of atomic and molecular pairs [6, 7], an approximation also independently developed by another group [8]. Fully consistent electrodynamics gave a perceptible, but not dramatic, statistical improvement [9] for those atomic and molecular dimers tested. However, one should use an improved approximation in cases where the fragment polarizabilities have significant anisotropy and also for the asymptotic interaction between two parallel surfaces, which is not given exactly by this approximation. In future versions of the functional, it will probably be possible to eliminate this S -expansion, keeping only the expansion in terms of the *difference* between ϵ and $1 - V\chi$. The retention of the anisotropic components of ϵ should also be possible.

In order to evaluate Eq. (4), we need a simple approximation for S , as a functional of the density. This choice is constrained by a number of exact relationships. In a plane-wave representation, $S_{\vec{q},\vec{q}'}$, one has the requirements (i) $S_{\vec{q},\vec{q}'}(\omega) \rightarrow -(4\pi e^2/m\omega^2)n_{\vec{q}-\vec{q}'}$ at large frequencies (the f -sum rule), where $n_{\vec{k}}$ is the Fourier transform of the density; (ii) $\int_{-\infty}^{\infty} du S_{\vec{q},\vec{q}'}(iu) \rightarrow 8\pi^2 Ne^2/q^2$ for large q , where N is the electron number, to reproduce the exactly known self-correlation; (iii) $S_{\vec{q},\vec{q}'} = S_{-\vec{q}',-\vec{q}}$ for time reversal invariance; (iv) a finite $S_{\vec{q},\vec{q}'}(\omega)$ for vanishing q or q' at all non-zero values of ω , to give an exchange-correlation hole with the correct volume (charge conservation).

An approximate S inspired by the plasmon-pole model successfully applied earlier [10] takes $S_{\vec{q},\vec{q}'} = \frac{1}{2}[\tilde{S}_{\vec{q},\vec{q}'} + \tilde{S}_{-\vec{q}',-\vec{q}}]$ where

$$\tilde{S}_{\vec{q},\vec{q}'} = \int d^3r e^{-i(\vec{q}-\vec{q}')\cdot\vec{r}} \frac{4\pi n(\vec{r})e^2/m}{(\omega + \omega_q(\vec{r}))(-\omega + \omega_{q'}(\vec{r}))}. \quad (5)$$

We will take $\omega_q(\vec{r})$ to be a function of the local density at point $\vec{r}$ and its gradient. The above S satisfies all the constraints provided that $\omega_q \rightarrow q^2/2m$ for large q .

To facilitate the numerical evaluation we choose an ω_q which depends on a single length scale l , and switches from its small- q form $\omega_q = 1/2ml^2$ to its large- q form (above) when $q \sim 1/l$. We make an arbitrary choice for the switching function letting $[1 - e^{-(ql)^2}]\omega_q = q^2/2m$. The quantity l will be a function of position, and for convenience later we define $q_0^2 = \gamma/l^2$, where $\gamma = 4\pi/9$. Hence, letting $h(y) = 1 - e^{-y^2}$, we may write

$$\omega_q(\vec{r}) = \frac{q^2}{2m} \frac{1}{h(q/q_0(\vec{r}))}. \quad (6)$$

It would of course be preferable to use a form with a second length scale, as would be provided by a linear

term $\propto qv_F$ in the ω_q dispersion [3, 4, 10], but this source of error is mitigated by a continuously variable choice of q_0 based on the density. We make this choice so the exchange-correlation energy density $\epsilon_{xc}^0(\vec{r})$ defined by

$$E_{xc}^0 = \int d^3r \epsilon_{xc}^0(\vec{r})n(\vec{r}) \quad (7)$$

produced by this choice of $q_0(\vec{r})$ corresponds to that of a full calculation.

The expression for E_{xc}^0 corresponding to the approximation (3) for E_c^{nl} is simply

$$E_{xc}^0 \approx \int_0^\infty \frac{du}{2\pi} \text{tr}(\ln \epsilon) - E_{\text{self}} \approx \int_0^\infty \frac{du}{2\pi} \text{tr} S - E_{\text{self}}, \quad (8)$$

where E_{self} subtracts off the internal Coulomb self-energy of each electron. As was done previously, we expand to lowest order in S in the second equality. Substituting for S using (5), integrating over $u = -i\omega$, and using Eq. (7) gives

$$\epsilon_{xc}^0(\vec{r}) = \frac{\pi e^2}{m} \int \frac{d^3q}{(2\pi)^3} \left[\frac{1}{\omega_q(\vec{r})} - \frac{2m}{q^2} \right], \quad (9)$$

where the second term in the brackets is the self-energy subtraction written explicitly. Upon substitution from (6), one finds

$$\epsilon_{xc}^0(\vec{r}) = \frac{e^2 q_0(\vec{r})}{\pi} \int_0^\infty dy [h(y) - 1] = -\frac{3e^2}{4\pi} q_0(\vec{r}). \quad (10)$$

Approximations for ϵ_{xc} are conveniently expressed as their ratio to the LDA exchange value $\epsilon_x^{\text{LDA}} = -3e^2 k_F/4\pi$, where $k_F^3 = 3\pi^2 n$. Equation (10) then implies that the local value of the parameter q_0 is simply given by the local value of k_F modulated by an easily understood energy ratio, that is

$$q_0(\vec{r}) = \frac{\epsilon_{xc}^0(\vec{r})}{\epsilon_x^{\text{LDA}}(\vec{r})} k_F(\vec{r}). \quad (11)$$

Equation (11) is used to determine the q_0 value to be used in Eq. (6), continuously as a function of position. For this purpose we use LDA with gradient corrections

$$\epsilon_{xc}^0 \approx \epsilon_{xc}^{\text{LDA}} - \epsilon_x^{\text{LDA}} \left[\frac{Z_{ab}}{9} \left(\frac{\nabla n}{2k_F n} \right)^2 \right], \quad (12)$$

where $Z_{ab} = -0.8491$. This is the contribution labeled “screened exchange” in the table in the review of Ref. 11, where the relationship to the original work of various authors, who obtained this quantity from first principles, is discussed. The remaining contribution Z_c , labeled “fluctuation” in Ref. 11, is part of E_c^{nl} , and hence inappropriate to include as part of E_{xc}^0 . It comes from a term in perturbation theory that gives a long range vdW-like interaction if taken at long range instead of in an expansion in gradients [12].

Writing Eq. (4) in a plane-wave representation gives

$$E_{xc}^{nl} = \int_0^\infty \frac{du}{4\pi} \sum_{\vec{q}, \vec{q}'} [1 - (\hat{q} \cdot \hat{q}')^2] S_{\vec{q}, \vec{q}'} S_{\vec{q}', \vec{q}}. \quad (13)$$

This may be straightforwardly, but tediously, expressed in the form of Eq. (1), where the kernel ϕ is given by

$$\begin{aligned} \phi(\vec{r}, \vec{r}') &= \frac{2me^4}{\pi^2} \int_0^\infty a^2 da \int_0^\infty b^2 db W(a, b) \\ &\times T(\nu(a), \nu(b), \nu'(a), \nu'(b)), \end{aligned} \quad (14)$$

where

$$\begin{aligned} T(w, x, y, z) &= \frac{1}{2} \left[\frac{1}{w+x} + \frac{1}{y+z} \right] \\ &\times \left[\frac{1}{(w+y)(x+z)} + \frac{1}{(w+z)(y+x)} \right] \end{aligned} \quad (15)$$

and

$$\begin{aligned} W(a, b) &= 2[(3 - a^2)b \cos b \sin a + (3 - b^2)a \cos a \sin b \\ &+ (a^2 + b^2 - 3) \sin a \sin b - 3ab \cos a \cos b] / a^3 b^3. \end{aligned} \quad (16)$$

The quantities ν and ν' are given by $\nu(y) = y^2/2h(y/d)$ and $\nu'(y) = y^2/2h(y/d')$, with $d = |\vec{r} - \vec{r}'|q_0(\vec{r})$ and $d' = |\vec{r} - \vec{r}'|q_0(\vec{r}')$, where q_0 is given by Eq. (11). The kernel ϕ thus depends on $\vec{r}$ and $\vec{r}'$ only through d and d' , so that ϕ can be tabulated in advance in terms of these two variables, or better yet in terms of the sum and difference variables D and δ defined by $d = D(1 + \delta)$ and $d' = D(1 - \delta)$. Then $0 \leq D < \infty$ and $0 \leq |\delta| < 1$. For large d and d' , the asymptotic form is

$$\phi \rightarrow -\frac{C}{d^2 d'^2 (d^2 + d'^2)}, \quad (17)$$

where $C = 12(4\pi/9)^3 me^4$. In Fig. 1 we show a plot of $4\pi D^2 \phi$ vs. D for several values of δ . The integral of the $\delta = 0$ curve vanishes as it must.

The numerical work uses Eq. (2) for the correlation functional coupled with the Zhang-Yang revPBE [13] exchange functional. This choice is motivated by the work of Wu *et al.* [14], which pointed out that a more standard GGA predicts substantial binding in rare gas dimers from exchange alone, a feature absent for exact Hartree-Fock exchange. We found [4] that revPBE exchange does not have this property, so by using it we insure that vdW binding, a correlation effect, actually comes from the correlation term in our approximation scheme [15].

In Table I our calculated values for the coefficient C_6 in the asymptotic interaction $-C_6/R^6$ between the elements of several dimers are compared with those from previous related calculations as well as reference values, which also have some uncertainty. The new values track those from Ref. 6 which used the same approximation for the electrostatics, and are only slightly inferior to the

values obtained in Ref. 9, where self-consistent electrostatics was used.

Figure 2 shows the calculated binding-energy curves as functions of separation for Ar and Kr dimers. The comparison with experimental values for the binding energy and distance [16] illustrate the promise of vdW-DF for such systems. The binding-energy curves for a benzene dimer in the atop-parallel configuration in Fig. 3 illustrate the importance of choosing the right GGA flavor, *i.e.* revPBE [13], to avoid erroneous attraction in exchange-only accounts, as discussed in Refs. 4 and 14. Figure 4 illustrates the relative agreement between modern wavefunction based calculations [17] and our vdW-DF method, and the importance of not using the GGA alone.

The above moderate successes of the vdW-DF proposed here suggest that its use, along with future improvements, may be a way to proceed for calculating properties of vdW bound molecules that are too large for wavefunction based methods to be useful.

Financial support from the Swedish Foundation for Strategic Research via Materials Consortia #9 and ATOMICS and the Swedish Scientific Council is gratefully acknowledged. Work by M.D. and D.C.L. was supported in part by NSF Grant DMR 00-93070.




FIG. 1: The kernel ϕ in Eq. (1) as a function of the dimensionless D parameter for several values of the asymmetry parameter δ , as defined in the text.

-
- [1] V. N. Staroverov, G. E. Scuseria, J. Tao, and J. P. Perdew, *J. Chem. Phys.* **119**, 12129 (2003).
 - [2] V. N. Staroverov, G. E. Scuseria, J. Tao, and J. P. Perdew, *Phys. Rev. B* **69**, 0351xxx (2004).
 - [3] H. Rydberg *et al.*, *Phys. Rev. Lett.* **91**, 126402 (2003).
 - [4] D. C. Langreth *et al.*, *Int. J. Quantum Chem.* **xx**, xxx (submitted), preprint: <http://www.physics.rutgers.edu/~langreth/dft2003.pdf>.




FIG. 2: Calculated interaction energy between two Ar atoms (dashed curves) and between two Kr atoms (solid curves). The experimental equilibrium data [16] are shown for comparison with the full vdW-DF, and the GGA predictions in the revPBE flavor are also shown.




FIG. 3: Interaction energy between two benzene molecules in the atop-parallel configuration as predicted by the exchange-only part of two GGA functionals. The full Hartree-Fock results [17] are shown for comparison.




FIG. 4: Full interaction energy between two benzene molecules in the atop-parallel configuration using the vdW-DF functional. For comparison, we show recent results [17] using coupled-cluster [CCSD(T)] and perturbation-theoretic (MP2) methods, as well as the prediction of two flavors of GGA. Our vdW-DF would have given an equilibrium separation closer to those from the wavefunction calculations, if exact HF exchange (see Fig. 3) had been used instead of revPBE exchange.

TABLE I: C_6 values for dimers (Rydberg atomic units). Present: from Eq. (17). Similar: calculations from Ref. 6 using the same electrostatics approximation. Unified: calculations from Ref. 9 using self-consistent electrostatics. Reference: sources cited in Ref. 9.

Dimer	Present	Similar	Unified	Reference
He	4.8	4	2.58	2.92
Ne	14.6	12	15.0	13.8
Ar	124	126	143	134
Kr	238	245	291	266
Xe	516	520	663	597
Mg	1598	1513	1230	1240

- [5] If both spatial arguments of $\tilde{\chi}$ are on one fragment, its value is still influenced by *spectator* interactions with other fragments.
- [6] Y. Andersson, D. C. Langreth, and B. I. Lundqvist, Phys. Rev. Lett. **76**, 102 (1996).
- [7] Y. Andersson and H. Rydberg, Physica Scripta **60**, 211 (1999).
- [8] J. F. Dobson and B. P. Dinte, Phys. Rev. Lett. **76**, 1780 (1996).
- [9] E. Hult, H. Rydberg, B. I. Lundqvist, and D. C. Langreth, Phys. Rev. B **59**, 4708 (1998).
- [10] H. Rydberg, B. I. Lundqvist, D. C. Langreth, and M. Dion, Phys. Rev. B **62**, 6997 (2000).
- [11] D. C. Langreth and S. H. Vosko, Advances in Quantum Chemistry **21**, 175 (1990).
- [12] Alternatively, an expansion of Eq. (4) in gradients gives

- the precise structure of Z_c , but with the triangular vertex parts [see Fig. 3(c) in Ref. 11] evaluated in local-density approximations. The gross failure of the old-fashioned gradient approximation, due to the unwitting attempt to expand the van der Waals interaction in terms of semilocal gradients, is obviated here by treating the van der Waals term in a more appropriate approximation.
- [13] Y. Zhang and W. Yang, Phys. Rev. Lett. **80**, 890 (1998).
- [14] X. Wu *et al.*, J. Chem. Phys. **115**, 8748 (2001).
- [15] The calculations were done with the DACAPO plane-wave pseudopotential code, using the full revPBE exchange-correlation potential. E_c^{nl} was added and the beyond LDA part of the revPBE correlation energy subtracted as post GGA perturbations. The core densities were restored by hand and found to make a negligible difference in E_c^{nl} .
- [16] J. Ogilvie and F. Wang, J. Mol. Struct. **273**, 277 (1992).
- [17] S. Tsuzuki, K. Honda, M. Mikami, and K. Tanabe, J. Am. Chem. Soc. **124**, 104 (2002), and references therein.